

pubs.acs.org/JPCA

#### Article

# Reaction Dynamics Study of the Molecular Hydrogen Loss Channel in the Elementary Reactions of Ground-State Silicon Atoms (Si(<sup>3</sup>P)) With 1- and 2-Methyl-1,3-Butadiene ( $C_5H_8$ )

Zhenghai Yang, Chao He, Shane Goettl, and Ralf I. Kaiser\*



suggests that the methyl group-although a spectator from the chemical viewpoint-can influence the disposal of the angular momentum into the rotational excitation of the final product.

# **1. INTRODUCTION**

Carbon (C) and silicon (Si) both belong to main group XIV and hence have each four valence electrons along with a  ${}^{3}P_{i}$ electronic ground state. Despite their isovalency, carbon and silicon depict distinct chemical dynamics upon reaction with unsaturated hydrocarbons (alkynes, alkenes, aromatics).<sup>1-11</sup> Reactions of  $C({}^{3}P_{i})$  with unsaturated hydrocarbons are very fast close to gas kinetic limits and hold rate constants of a few  $10^{-10}$  cm<sup>3</sup> s<sup>-1</sup> over temperatures from 15 to 300 K.<sup>12–17</sup> These processes are barrierless and proceed via addition of the carbon atom to one and/or both carbon atoms of the unsaturated bond; the initial reaction intermediates either isomerize through ring opening/ring closure and facile hydrogen shifts eventually forming highly unsaturated, often resonantly stabilized free radicals (RSFRs) through the carbon versus hydrogen atom exchange pathways in overall exoergic reactions.<sup>18-22</sup> These RSFRs are frequently substituted propargyl radicals as derived from the replacement of a hydrogen atom of propargyl (1) by methyl  $(CH_3)$ , ethynyl (CCH), or vinyl  $(C_2H_3)$  groups and might act as a feedstock of (doubly) substituted benzenes in combustion flames and in deep space. With the exception of the carbon-acetylene system, 5,8,11,23 which also reveals nonadiabatic reaction dynamics through intersystem crossing (ISC) from the triplet to the singlet surface accompanied by molecular hydrogen loss

and tricarbon  $(C_3)$  formation, all carbon-hydrocarbon exchange reactions proceed likely on the triplet surface.

On the other hand, the reactivities of  $Si({}^{3}P_{i})$  with hydrocarbon systems are quite distinct. Upon collisions with C1-C3 hydrocarbons, silicon does not react at all under single collision conditions through exoergic reactions. A crossed molecular beam study probed the Si(3P) plus acetylene  $(C_2H_2)$  system, revealing the formation of the linear ethynylsilylidyne (SiCCH) molecule and atomic hydrogen at a collision energy of 101.6  $\pm$  1.6 kJ mol<sup>-1</sup> and a reaction endoergicity of +84  $\pm$  6 kJ mol<sup>-1.24</sup> No reactive scattering signal could be detected for experiments under single collision conditions of Si(<sup>3</sup>P) with any hydrocarbon up to C3 including the C<sub>3</sub>H<sub>4</sub> isomers methylacetylene (CH<sub>3</sub>CCH) and allene  $(CH_2CCH_2)$  at collision energies up to 30 kJ mol<sup>-1</sup>, indicating that endoergicities and/or reaction barriers in this reaction system are insurmountable. This lack of reactivity seems to be in contrast to early kinetics studies by Canosa et al.,<sup>25</sup> Basu and Husain,<sup>26,27</sup> which suggest that the reactions of Si(<sup>3</sup>P) with

Received: April 3, 2021 Revised: May 26, 2021 Published: June 7, 2021



## The Journal of Physical Chemistry A

unsaturated hydrocarbons are close to gas kinetics values with rate constants of a few  $10^{-10}$  cm<sup>3</sup> s<sup>-1</sup>. However, only the decay kinetics of the silicon atom were probed in these studies, without the identification of the reaction products. In fact, these reaction products are of critical importance since they can help unraveling fundamental constraints on the exotic chemical bonding and reaction mechanism(s) (bimolecular versus three-body reactions) of organosilicon molecules. This discrepancy might be explained through the formation of collision complexes between silicon and the hydrocarbons followed by rapid collisional stabilization. The results of calculations by Talbi also suggested that the rapid rate constants for the silicon plus acetylene and silicon plus ethylene reactions were due to the intervention of a third body. Within the hypotheses of efficient intersystem crossing to the singlet surface, the barrierless formation of some singlet and triplet SiC<sub>2</sub>H<sub>2</sub> and SiC<sub>2</sub>H<sub>4</sub> isomers should be inhibited.<sup>28</sup> Bimolecular reactions of ground-state oxygen atoms  $O(^{3}P)$ with unsaturated hydrocarbons like acetylene,<sup>29</sup> ethylene,<sup>30</sup> propene,<sup>31</sup> allene,<sup>32</sup> and 1-butene<sup>33</sup> have been studied previously. These multichannel reactions involve important nonadiabatic effects, and products can be formed through exoergic reactions even in reactions with C1-C3 hydrocarbons; this finding is quite different from the silicon reactions. Very recently, crossed beam experiments revealed that Si(<sup>3</sup>P) reacts excergically with four  $C_4H_6$  isomers, i.e., 2butyne (CH<sub>3</sub>CCCH<sub>3</sub>),<sup>34</sup> 1,3-butadiene (H<sub>2</sub>CCHCHCH<sub>2</sub>), 1,2-butadiene (CH2CCHCH3), and 1-butyne (CH<sub>3</sub>CH<sub>2</sub>CCH), producing multiple singlet  $SiC_4H_4$  isomers such as 2-methylene-1-silacyclobutene (SiC<sub>2</sub>H<sub>2</sub>CCH<sub>2</sub>; X<sup>1</sup>A) along with molecular hydrogen  $(H_2)$  through nonadiabatic reaction dynamics.<sup>35</sup> Considering the lack of rapid reaction of C1 to C3 hydrocarbons with ground-state atomic silicon, but the facile nonadiabatic dynamics of atomic silicon with the four  $C_4H_6$  isomers, the chemical dynamics of Si(<sup>3</sup>P<sub>i</sub>) with hydrocarbons still remain poorly understood, and a more systematic investigation is required.

Here, the experimental results of the Si(<sup>3</sup>P) reactions with two  $C_5H_8$  isomers,1-methyl-1,3-butadiene and 2-methyl-1,3butadiene, are reported to expand our knowledge of the fundamental reaction mechanisms of the silicon–carbon chemistry in the gas phase. The formal linkage of these hydrocarbons to 1,3-butadiene can be explored by replacing one of the hydrogen atoms at the C1/C4 and C2/C3, respectively, by a methyl group. An understanding of the inherent (nonadiabatic) reaction dynamics will explore to what extend the methyl group simply acts as a spectator, induces sterical effects, and/or is actively engaged in the gas-phase chemistry such as via methane (CH<sub>4</sub>) elimination accompanied by formation of SiC<sub>4</sub>H<sub>4</sub> isomers.

#### 2. EXPERIMENTAL METHODS

The reactions of atomic silicon (Si; <sup>3</sup>P) with 1-methyl-1,3butadiene ( $\geq$ 97%; TCI America) and 2-methyl-1,3-butadiene (isoprene;  $\geq$ 99%; TCI America) were carried out in a crossed molecular beams machine.<sup>36–38</sup> Briefly, a supersonic atomic beam of ground-state silicon atoms was prepared in the primary source chamber in situ through ablation of a rotating silicon rod exploiting the 266 nm pulse (4th harmonic of a Nd:YAG; 5 ± 1 mJ per pulse; 30 Hz)<sup>39–42</sup> and entraining the ablated atoms in a pulsed neon (Ne; 99.999%; Airgas; 60 Hz; 4 atm) beam. The laser-induced fluorescence (LIF) spectra measured in the ablation source are presented in Figure 1; at



pubs.acs.org/JPCA

Figure 1. Laser-induced fluorescence (LIF) spectra of the (a) ground  $(^{3}P)$  and (b) excited  $(^{1}D)$  states of atomic silicon. The arrow in panel (b) indicates the expected position for the noted transition.

these experimental conditions, there are no electronically excited silicon atoms Si(1D) in the molecular beam. Thirty Hertz of ablation laser versus 60 Hz of the gas will allow a "laser on" minus "laser off" background subtraction. The neonseeded primary silicon beam was skimmed and then velocityselected by a 120 Hz high-precision four-slit chopper wheel (2057S024B, MC 5005 S RS, Faulhaber), which resulted in well-defined speed ratios (S) of 6.2  $\pm$  0.4 and 6.1  $\pm$  0.4 along with peak velocities ( $v_p$ ) of 975 ± 30 and 982 ± 30 m s<sup>-1</sup> (Table 1). As verified by the laser-induced fluorescence spectra, all silicon atoms of our neon-seeded silicon beam are in their electronic ground state (<sup>3</sup>P).<sup>43</sup> As for the secondary source, the hydrocarbon reactant 1-methyl-1,3-butadiene or 2methyl-1,3-butadiene was kept in a stainless-steel reservoir at 293 K under backing pressure of 400 or 430 Torr. The corresponding peak velocities and speed ratios of  $v_p = 711 \pm$ 20 m s<sup>-1</sup> with  $\tilde{S} = 8.5 \pm 0.7$  and  $v_p = 721 \pm 20$  with  $\tilde{S} = 8.5 \pm$ 0.6 were derived, respectively. The hydrocarbon beams crossed perpendicularly with the primary Si(<sup>3</sup>P) beam in the interaction region yielding center-of-mass (CM) angles of  $60.5 \pm 0.6$  and  $60.7 \pm 0.5^{\circ}$  and collision energies of  $14.4 \pm 0.4$ and 14.7  $\pm$  0.4 kJ mol<sup>-1</sup> (Table 1).

Table 1. Peak Velocity ( $\nu_p$ ) and Speed Ratios (S) of the Silicon (Si), 1,3-Pentadiene (CH<sub>2</sub>CHCHCHCH<sub>3</sub>), and Isoprene (CH<sub>2</sub>CHCCH<sub>3</sub>CH<sub>2</sub>) Beams along With the Corresponding Collision Energy ( $E_C$ ), Spread Ratio of Collision Energy (S), and Center-of-Mass Angle ( $\Theta_{CM}$ ) for Each Reactive Scattering Experiment

pubs.acs.org/JPCA

beam	$\nu_{\rm p}~({\rm m~s^{-1}})$	S	$E_{\rm c}$ (kJ mol <sup>-1</sup> )	S	$\Theta_{ m CM}$ (°)
Si ( <sup>3</sup> P)	$975 \pm 30$	$6.2 \pm 0.4$			
$CH_2CHCHCHCH_3$ (X <sup>1</sup> A')	$711 \pm 20$	$8.5 \pm 0.7$	$14.4 \pm 0.4$	$4.0 \pm 0.3$	$60.5 \pm 0.6$
Si ( <sup>3</sup> P)	$982 \pm 30$	$6.1 \pm 0.4$			
$CH_2CHCCH_3CH_2$ (X <sup>1</sup> A')	$721 \pm 20$	8.5 ± 0.6	$14.7 \pm 0.4$	$4.0 \pm 0.3$	$60.7 \pm 0.5$

In our experiment, a quadrupole mass filter was exploited to mass filter the ionized (80 eV, 2 mA) reactive scattering products, a 1.2 MHz oscillator, and a Daly-type time-of-flight (TOF) detector, which is installed in a ultrahigh vacuum  $(<10^{-11} \text{ Torr})$  chamber, were also used. The recorded TOF spectra eventually provide the product angular distribution in the laboratory frame and provide the basis for the transformation of the laboratory data into the center-of-mass (CM) frame through the use of a forward-convolution routine. This results in the extraction of the CM angular flux distribution  $T(\theta)$  and translational energy flux distribution  $P(E_{\rm T})$ . Finally, the best fits of the lab angular distribution and TOF spectra are derived by optimizing the  $T(\theta)$  and  $P(E_{\rm T})$  functions. The reactive differential cross section can be displayed via the flux contour map with the CM velocity u and scattering angle  $\theta$ , i.e.,  $I(u, \theta) \sim P(u) \times T(\theta)$ .

## 3. RESULTS

3.1. Laboratory Frame. Our scattering experiments explored signal at mass-to-charge ratios (m/z) 98 to 94 and 82 to 80 to investigate the presence of molecular  $(H_2)$  and atomic (H) hydrogen loss channels, methylene (CH<sub>2</sub>), methyl (CH<sub>3</sub>), and methane (CH<sub>4</sub>) elimination pathways along with adducts. Accounting for the natural abundances of three silicon isotopes (<sup>30</sup>Si (3.1%), <sup>29</sup>Si (4.7%), and <sup>28</sup>Si (92.2%)) as well as carbon (<sup>13</sup>C (1.1%), <sup>12</sup>C (98.9%)), we were not able to detect signal at m/z 98 and 97. Signal was observed at m/z 96 signal at m/2 98 and 97. Signal was observed at m/2 96 ( ${}^{30}\text{SiC}_{5}\text{H}_{6}^{+}$ ,  ${}^{30}\text{Si}^{13}\text{CC}_{4}\text{H}_{5}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{7}^{+}$ ,  ${}^{29}\text{Si}^{13}\text{CC}_{4}\text{H}_{6}^{+}$ ,  ${}^{28}\text{Si}^{13}\text{CC}_{4}\text{H}_{7}^{+}$ ,  ${}^{28}\text{SiC}_{5}\text{H}_{8}^{+}$ ), 95 ( ${}^{30}\text{SiC}_{5}\text{H}_{5}^{+}$ ,  ${}^{30}\text{Si}^{13}\text{CC}_{4}\text{H}_{4}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{6}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{6}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{6}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{7}^{+}$ ), and 94 ( ${}^{30}\text{SiC}_{5}\text{H}_{4}^{+}$ ,  ${}^{30}\text{Si}^{13}\text{CC}_{4}\text{H}_{3}^{+}$ ,  ${}^{29}\text{SiC}_{5}\text{H}_{5}^{+}$ ,  ${}^{29}\text{Si}^{13}\text{CC}_{4}\text{H}_{4}^{+}$ ,  ${}^{28}\text{Si}^{13}\text{CC}_{4}\text{H}_{5}^{+}$ , and  ${}^{28}\text{SiC}_{5}\text{H}_{6}^{+}$ ). Since after scaling, the TOFs at m/z = 96, 95, and 94 reveal identical patterns and hence do overlap, a single reaction channel is likely. These findings propose the absence of adducts  $({}^{30}\text{SiC}_5\text{H}_8^+/{}^{29}\text{SiC}_5\text{H}_8^+/{}^{28}\text{SiC}_5\text{H}_8^+)$  and the nonexistence of the atomic hydrogen loss channel  $({}^{30}\text{SiC}_5\text{H}_7^+/{}^{29}\text{SiC}_5\text{H}_7^+/{}^{28}\text{SiC}_5\text{H}_7^+)$ . Quantitatively spoken, signal at m/z 96 to 95 was observed at levels of close to 4% (4  $\pm$ 1% for 1,3-pentadiene,  $3 \pm 1\%$  for isoprene), 10% (12  $\pm 3\%$ for 1,3-pentadiene,  $11 \pm 2\%$  for isoprene) compared to m/z =94, which is close to the calculated isotopic ratios. Signal at m/z 96, 95, and 94 can be attributed to  ${}^{30}SiC_5H_6^+$ ,  ${}^{29}SiC_5H_6^+$ ,  ${}^{28}\text{Si}{}^{13}\text{CC}_4\text{H}_6^+$ , and  ${}^{28}\text{SiC}_5\text{H}_6^+$  (hereafter: SiC<sub>5</sub>H<sub>6</sub>), respectively, along with their fragment ions  $({}^{30}SiC_5H_5{}^+/{}^{30}SiC_5H_4{}^+/{}^{29}SiC_5H_5{}^+/{}^{28}Si^{13}CC_4H_5{}^+)$  (Figures 2a and 2b). Therefore, according to the laboratory data, there is strong evidence of the molecular hydrogen loss pathway, leading to the formation of SiC5H6 isomers in the reaction of  $Si(^{3}P)$  with both C<sub>5</sub>H<sub>8</sub> isomers, 1,3-pentadiene and isoprene. For both systems, TOF spectra for the molecular hydrogen loss channel were collected at m/z = 94, i.e., the dominant ion, in steps of 2.5°. The TOF spectra are very broad from at least



Article

**Figure 2a.** Laboratory angular distribution (a) and time-of-flight spectra (b) recorded in the reaction of the ground-state atomic silicon with 1,3-pentadiene at a mass-to-charge ratio of m/z = 94. The normalized experimental distribution is represented by the solid circles with  $\pm 1\sigma$  error bars and the experimental time-of-flight spectra data points are indicated by the open circles. The best fits represented by the red lines are derived from the optimized center-of-mass (CM) functions, as depicted in Figure 3a. Silicon, carbon, and hydrogen are color-coded in purple, gray, and white, respectively.

350 to 900  $\mu$ s and peak between 400 and 500  $\mu$ s. The normalized laboratory angular distributions span at least 35° and reveal maxima close to their respective CM angles (Figures 2a and 2b). Their symmetries around the CM angles propose indirect reaction mechanisms with at least one SiC<sub>5</sub>H<sub>8</sub> reaction intermediate and molecular hydrogen (H<sub>2</sub>) loss to form the SiC<sub>5</sub>H<sub>6</sub> isomer(s).

**3.2.** Center-of-Mass Frame. An  $E_{\rm C}^{-1/3}$  dependence reaction cross section for a dipole-induced dipole interaction and a single reaction channel fit with the mass combination Si (28 amu) + C<sub>5</sub>H<sub>8</sub> (68 amu)  $\rightarrow$  SiC<sub>5</sub>H<sub>6</sub> (94 amu) + H<sub>2</sub> (2 amu) were utilized for both systems.<sup>44</sup> The best-fitting CM functions are compiled in red lines in Figures 3a and 3b and 1 $\sigma$  error limits of the scattering experiment are represented by the hatched areas of the  $P(E_{\rm T})$  and  $T(\theta)$  functions. First, let us consider the center-of-mass translational energy distributions. The maximum energy  $E_{\rm max}$  available to the translation degrees of freedom of the nascent SiC<sub>5</sub>H<sub>6</sub> plus molecular hydrogen



**Figure 2b.** Laboratory angular distribution (a) and time-of-flight spectra (b) recorded in the reaction of the ground-state atomic silicon with isoprene at a mass-to-charge ratio of m/z = 94. The normalized experimental distribution is represented by the solid circles with  $\pm 1\sigma$  error bars and the experimental time-of-flight spectra data points are indicated by the open circles. The best fits represented by the red lines are derived from the optimized center-of-mass (CM) functions, as depicted in Figure 3b. Silicon, carbon, and hydrogen are color-coded in purple, gray, and white, respectively.

products is determined as  $E_{\text{max}} = E_{\text{C}} - \Delta_{\text{r}}G$  through energy conservation by summing the reaction energy and the collision energy. Accordingly, by subtracting the collision energy from the maximum translational energy (Figures 3a and 3b), the reactions of Si(<sup>3</sup>P) with 1-methyl-1,3-butadiene and 2-methyl-1,3-butadiene to form SiC5H6 plus molecular hydrogen are exoergic by  $113 \pm 19$  and  $101 \pm 21$  kJ mol<sup>-1</sup>, respectively. The  $P(E_{T})$ s of the two systems peaked at translational energies of  $20 \pm 4$  and  $30 \pm 5$  kJ mol<sup>-1</sup>. These relatively high energies provide evidence of the existence of tight exit transition states in the molecular hydrogen elimination. Overall, both  $P(E_T)$ s are very similar in shape, albeit the silicon-1-methyl-1,3butadiene system revealing a slightly higher translational energy release. Comparable  $T(\theta)$  functions are also evident for both systems with best-fitting  $T(\theta)$  achieved for a product flux over the entire angular range. The forward-backward symmetries and sideways scattering (maximum at  $90^\circ$ ) suggest the presence of a decomposing  $SiC_5H_8$  intermediate (indirect scattering dynamics) with a lifetime comparable to or exceeding its rotational period;<sup>45,46</sup> the sideways scattering reveals significant geometrical constraints and suggests that molecular hydrogen elimination occurs nearly parallel to the total angular momentum vector and hence perpendicularly to the rotational plane of the decomposing intermediate.<sup>4</sup> However, the more pronounced peaking in the silicon-1methyl-1,3-butadiene system is evident (Figure 3a).

#### 4. DISCUSSION

Our experimental results reveal that atomic silicon reacts exoergically with 1-methyl- and 2-methyl-1,3-butadiene to  $SiC_{5}H_{6}$  organosilicon product(s) together with molecular hydrogen (H<sub>2</sub>) through tight exit transition states and longlived  $SiC_{5}H_{8}$  reaction intermediate(s). The molecular formula of these newly synthesized organosilicon reaction products has



**Figure 3a.** Center-of-mass (CM) translational energy flux distribution (a), angular distribution (b), and the corresponding flux contour map (c) for the reaction of ground-state atomic silicon with 1,3-pentadiene. The  $1\sigma$  error limits of the scattering experiment are represented by the shaded areas and the best-fit functions are defined by the red solid lines. The flux contour map symbolizes the reactive scattering products as a function of the CM scattering angle ( $\theta$ ) and product velocity (u). The flux gradient from high (H) intensity to low (L) intensity is indicated by the color bar. Silicon, carbon, and hydrogen are color-coded in purple, gray, and white, respectively.

pubs.acs.org/JPCA



**Figure 3b.** Center-of-Mass (CM) translational energy flux distribution (a), angular distribution (b) and the corresponding flux contour map (c) for the reaction of ground-state atomic silicon with isoprene. The  $1\sigma$  error limits of the scattering experiment are represented by the shaded areas and the best-fit functions are defined by red solid lines. The flux contour map symbolizes the reactive scattering products as a function of the CM scattering angle ( $\theta$ ) and product velocity (u). The flux gradient from high (H) intensity to low (L) intensity is indicated by the color bar. Silicon, carbon, and hydrogen are color-coded in purple, gray, and white, respectively.



**Figure 4a.** Inferred reaction pathway of ground-state silicon atom  $(Si(^{3}P))$  with 1,3-pentadiene  $(CH_2CHCHCHCH_3)$  by analogy with the 1,3-butadiene case (Figure 8 in ref 35), leading via molecular hydrogen loss to methyl-substituted c-SiCHCHCCH<sub>2</sub> products. Two proposed pathways accounting for the barrierless addition of the silicon atom to the different carbon–carbon double bond of the 1,3-pentadiene are color-coded in black and red, respectively.

pubs.acs.org/JPCA

Article



Figure 4b. Inferred reaction pathway of ground-state silicon atom  $(Si(^{3}P))$  with isoprene  $(CH_{2}CHCCH_{3}CH_{2})$  by analogy with the 1,3-butadiene case (Figure 8 in ref 35) leading via molecular hydrogen loss to methyl-substituted c-SiCHCHCCH<sub>2</sub> products.

been determined  $(SiC_5H_6)$ ; our study will also propose the molecular structure, inherent formation mechanism(s), and reaction dynamics of the product isomer(s). Traditionally, feasible reaction products are determined by comparing the experimentally derived reaction energies for each system  $(-113 \pm 19 \text{ and } -101 \pm 21 \text{ kJ mol}^{-1})$  with reaction energies obtained from electronic structure calculations.<sup>34,35</sup> However, a full theoretical analysis of the SiC5H8 potential energy surface (PES) and conical intersections along with a statistical Rice-Ramsperger-Kassel-Marcus (RRKM) treatment is well beyond the scope of this dissemination considering the complexity of both systems. Considering the fact that we only observed molecular hydrogen loss, but neither atomic hydrogen, methyl, nor methane loss pathways, we are attempting an elucidation of the reaction dynamics by comparing the reaction dynamics of the Si-1,3-butadiene reaction with the methyl-substituted systems obtained here. Briefly, the reaction of ground-state silicon atoms with 1,3butadiene is initiated by a barrierless addition of the silicon atom to one of the carbon-carbon double bonds, forming a triplet reaction intermediate <sup>3</sup>i1 stabilized by 91 kJ mol<sup>-1</sup> (Figure 8 in ref 35). For the energetically most favorable channel, the latter can undergo facile intersystem crossing to the singlet manifold, which leads to an il intermediate, which resides in a 190 kJ mol<sup>-1</sup> potential energy well. An atomic hydrogen migration from the H<sub>2</sub>C moiety within the threemembered ring is accompanied by ring opening to i2 followed by two trans-cis isomerization (i3, i4). The latter eventually emits molecular hydrogen followed by cyclization to 2methylene-1-silacyclobutene (c-SiCHCHCCH<sub>2</sub>); this mechanism reveals that the hydrogen atoms in molecular hydrogen originate from the C1 and C3 carbon atoms each in an overall exoergic reaction  $(-120 \pm 5 \text{ kJ mol}^{-1})$ . These dynamics correlate with the results from the reaction of ground-state silicon with 1,1,4,4-D<sub>4</sub>- and 2,3-D<sub>2</sub>-1,3-butadiene and the identification of the HD channel. Let us evaluate what happens if this mechanism is exported to the 1-methyl- and 2-methyl-1,3-butadiene reactants (Figures 4a and 4b). Note that due to the methyl group, the carbon-carbon double bonds are not chemically equivalent as they are in 1,3-butadiene. These pathways could lead to three triplet collision complexes (<sup>3</sup>i5, <sup>3</sup>i9, and <sup>3</sup>i13), which could undergo intersystem crossing to singlet i5, i9, and i13. Successive hydrogen shifts accompanied by ring openings (i6, i10, and i14) followed by trans/cis isomerization eventually led to i8, i12, and i16, which could emit molecular hydrogen to yield methyl-substituted analog structures of 2-methylene-1-silacyclobutene (c-SiCHCHCCH<sub>2</sub>). From the chemical viewpoint, the methyl group may act as a spectator in both reactions. However, considering the distinct structures and moment of inertia of the decomposing complexes i8, i12, and i16, from the dynamical viewpoint, the methyl group influences the chemical dynamics, leading to a more pronounced sideways scattering in the silicon-1-methyl-1,3-butadiene system compared to the reaction with 2-methyl-1,3-butadiene (isoprene). It shall be noted that the proposed dynamics represent one of possible multiple (less exoergic) additional reaction pathways, which remain to be explored.

## The Journal of Physical Chemistry A

## 5. CONCLUSIONS

In conclusion, bimolecular gas-phase reactions of ground-state atomic silicon (Si; <sup>3</sup>P) with 1- and 2-methyl-1,3-butadiene were studied; the reactions revealed indirect scattering dynamics through long-lived SiC<sub>5</sub>H<sub>8</sub> collision complex(es) together with a facile formation of  $SiC_sH_6$  isomer(s) along with molecular hydrogen. We propose-in an analogy of the derived center-of-mass functions of the silicon-1,3-butadiene system-that the reaction proceeds via nonadiabatic reaction dynamics through intersystem crossing (ISC) from the triplet to the singlet surface, yielding eventually to methyl-substituted 2-methylene-1-silacyclobutene isomers in overall exoergic reactions  $(-113 \pm 19 \text{ and } -101 \pm 21 \text{ kJ mol}^{-1})$  through tight exit transition states. Therefore, we may conclude that at least singly, alkyl-substituted 1,3-butadiene molecules react with atomic silicon to form eventually alkyl-substituted, cyclic 2-methylene-1-silacyclobutene products. Although the methyl group acts as a spectator from the chemical viewpoint, it does influence the disposal of angular momentum into the rotational excitation of the final products, as evident from distinct centerof-mass angular distributions of both systems explored here.

## AUTHOR INFORMATION

#### **Corresponding Author**

Ralf I. Kaiser – Department of Chemistry, University of Hawai'i at Manoa, Honolulu, Hawaii 96822, United States; orcid.org/0000-0002-7233-7206; Email: ralfk@ hawaii.edu

#### Authors

Zhenghai Yang – Department of Chemistry, University of Hawai'i at Manoa, Honolulu, Hawaii 96822, United States Chao He – Department of Chemistry, University of Hawai'i at

Manoa, Honolulu, Hawaii 96822, United States Shane Goettl – Department of Chemistry, University of

Hawai'i at Manoa, Honolulu, Hawaii 96822, United States

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.jpca.1c03023

#### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work at the University of Hawaii was supported by the US National Science Foundation (CHE-1853541).

#### REFERENCES

(1) Kaiser, R. I.; Stranges, D.; Bevsek, H. M.; Lee, Y. T.; Suits, A. G. Crossed-Beam Reaction of Carbon Atoms with Hydrocarbon Molecules. IV. Chemical Dynamics of Methylpropargyl Radical Formation,  $C_4H_5$ , from Reaction of  $C({}^{3}P_{j})$  with Propylene,  $C_3H_6(X^{1}A')$ . J. Chem. Phys. **1997**, 106, 4945–4953.

(2) Kaiser, R. I.; Sun, W.; Suits, A. G.; Lee, Y. T. Crossed Beam Reaction of Atomic Carbon,  $C({}^{3}P_{j})$ , with the Propargyl Radical,  $C_{3}H_{3}(X^{2}B_{2})$ : Observation of Diacetylene,  $C_{4}H_{2}(X^{1}\sum_{g}^{+})$ . J. Chem. Phys. **1997**, 107, 8713–8716.

(3) Hahndorf, I.; Lee, H. Y.; Mebel, A. M.; Lin, S. H.; Lee, Y. T.; Kaiser, R. I. A Combined Crossed Beam and Ab Initio Investigation on the Reaction of Carbon Species with  $C_4H_6$  isomers. I. The 1, 3-Butadiene Molecule,  $H_2CCHCHCH_2$  (X<sup>1</sup>A'). J. Chem. Phys. 2000, 113, 9622–9636.

(4) Hahndorf, I.; Lee, Y. T.; Kaiser, R. I.; Vereecken, L.; Peeters, J.; Bettinger, H. F.; Schreiner, P. R.; Schleyer, P. vR.; Allen, W. D.; Schaefer, H. F., III A Combined Crossed-Beam, Ab Initio, and RiceRamsperger–Kassel–Marcus Investigation of the Reaction of Carbon Atoms  $C({}^{3}P_{j})$  with Benzene,  $C_{6}H_{6}(X^{1}A_{1g})$  and  $D_{6}$ -Benzene,  $C_{6}D_{6}(X^{1}A_{1g})$ . J. Chem. Phys. **2002**, 116, 3248–3262.

(5) Gu, X.; Guo, Y.; Zhang, F.; Kaiser, R. I. Investigating the Chemical Dynamics of the Reaction of Ground-State Carbon Atoms with Acetylene and its Isotopomers. *J. Phys. Chem. A* 2007, 111, 2980–2992.

(6) Wilson, A. V.; Parker, D. S. N.; Zhang, F.; Kaiser, R. I. Crossed Beam Study of the Atom-Radical Reaction of Ground State Carbon Atoms ( $C(^{3}P)$ ) with the Vinyl Radical ( $C_{2}H_{3}(X^{2}A')$ ). *Phys. Chem. Chem. Phys.* **2012**, *14*, 477–481.

(7) Lucas, M.; Thomas, A. M.; Zhao, L.; Kaiser, R. I.; Kim, G. S.; Mebel, A. M. Gas-Phase Synthesis of the Elusive Cyclooctatetraenyl Radical ( $C_8H_7$ ) via Triplet Aromatic Cyclooctatetraene ( $C_8H_8$ ) and Non-Aromatic Cyclooctatriene ( $C_8H_8$ ) Intermediates. *Angew. Chem.* **2017**, *129*, 13843–13848.

(8) Leonori, F.; Petrucci, R.; Segoloni, E.; Bergeat, A.; Hickson, K. M.; Balucani, N.; Casavecchia, P. Unraveling the Dynamics of the  $C({}^{3}P, {}^{1}D) + C_{2}H_{2}$  Reactions by the Crossed Molecular Beam Scattering Technique. *J. Phys. Chem. A* **2008**, *112*, 1363–1379.

(9) Costes, M.; Daugey, N.; Naulin, C.; Bergeat, A.; Leonori, F.; Segoloni, E.; Petrucci, R.; Balucani, N.; Casavecchia, P. Crossed-Beam Studies on the Dynamics of the C +  $C_2H_2$  Interstellar Reaction Leading to Linear and Cyclic  $C_3H + H$  and  $C_3 + H_2$ . *Faraday Discuss.* **2006**, 133, 157–176.

(10) Geppert, W. D.; Naulin, C.; Costes, M.; Capozza, G.; Cartechini, L.; Casavecchia, P.; Gualberto Volpi, G. Combined Crossed-Beam Studies of  $C({}^{3}P_{J}) + C_{2}H_{4} \rightarrow C_{3}H_{3} + H$  Reaction Dynamics Between 0.49 and 30.8 kJ mol<sup>-1</sup>. *J. Chem. Phys.* **2003**, *119*, 10607–10617.

(11) Cartechini, L.; Bergeat, A.; Capozza, G.; Casavecchia, P.; Volpi, G. G.; Geppert, W. D.; Naulin, C.; Costes, M. Dynamics of the C +  $C_2H_2$  Reaction from Differential and Integral Cross-Section Measurements in Crossed-Beam Experiments. *J. Chem. Phys.* **2002**, *116*, 5603–5611.

(12) Chastaing, D.; Le Picard, S. D.; Sims, I. R.; Smith, I. W. M. Rate Coefficients for the Reactions of  $C({}^{3}P_{J})$  Atoms with  $C_{2}H_{2}$ ,  $C_{2}H_{4}$ , CH<sub>3</sub>CCH and H<sub>2</sub>CCCH<sub>2</sub> at Temperatures Down to 15 K. Astron. Astrophys. **2001**, 365, 241–247.

(13) Chastaing, D.; Le Picard, S. D.; Sims, I. R.; Smith, I. W. M.; Geppert, W. D.; Naulin, C.; Costes, M. Rate Coefficients and Cross-Sections for the Reactions of  $C({}^{3}P_{J})$  Atoms with Methylacetylene and Allene. *Chem. Phys. Lett.* **2000**, 331, 170–176.

(14) Chastaing, D.; James, P. L.; Sims, I. R.; Smith, I. W. M. Neutral–Neutral Reactions at the Temperatures of Interstellar Clouds: Rate Coefficients for Reactions of Atomic Carbon,  $C({}^{3}P)$ , with  $O_2$ ,  $C_2H_2$ ,  $C_2H_4$  and  $C_3H_6$  Down to 15 K. *Phys. Chem. Chem. Phys.* **1999**, *1*, 2247–2256.

(15) Haider, N.; Husain, D. Absolute Rate Data for the Reactions of Ground-State Atomic Carbon,  $C[2p^2(^3P_J)]$ , with Alkenes Investigated by Time-Resolved Atomic Resonance Absorption Spectroscopy in the Vacuum Ultraviolet. *J. Chem. Soc., Faraday Trans.* **1993**, *89*, 7–14.

(16) Haider, N.; Husain, D. The Collisional Behaviour of Ground State Atomic Carbon,  $C2p^{2}({}^{3}P_{J})$ , with Ethylene and Acetylene Investigated by Time-Resolved Atomic Resonance Absorption Spectroscopy in the Vacuum Ultraviolet. *J. Photochem. Photobiol., A* **1993**, 70, 119–124.

(17) Clary, D. C.; Haider, N.; Husain, D.; Kabir, M. Interstellar Carbon Chemistry: Reaction Rates of Neutral Atomic Carbon with Organic Molecules. *Astrophys. J.* **1994**, *422*, 416–422.

(18) Gu, X.; Kaiser, R. I.; Mebel, A. M. Chemistry of Energetically Activated Cumulenes—from Allene ( $H_2CCCH_2$ ) to Hexapentaene ( $H_2CCCCCH_2$ ). *ChemPhysChem* **2008**, *9*, 350–369.

(19) Kaiser, R. I.; Mebel, A. M. The Reactivity of Ground-State Carbon Atoms with Unsaturated Hydrocarbons in Combustion Flames and in the Interstellar Medium. *Int. Rev. Phys. Chem.* **2002**, *21*, 307–356.

#### The Journal of Physical Chemistry A

pubs.acs.org/JPCA

(20) Kaiser, R. I. Experimental Investigation on the Formation of Carbon-Bearing Molecules in the Interstellar Medium via Neutral–Neutral Reactions. *Chem. Rev.* **2002**, *102*, 1309–1358.

(21) Casavecchia, P.; Balucani, N.; Cartechini, L.; Capozza, G.; Bergeat, A.; Volpi, G. G. Crossed Beam Studies of Elementary Reactions of N and C Atoms and CN Radicals of Importance in Combustion. *Faraday Discuss.* **2001**, *119*, 27–49.

(22) Mebel, A. M.; Kaiser, R. I. Formation of Resonantly Stabilised Free Radicals via the Reactions of Atomic Carbon, Dicarbon, and Tricarbon with Unsaturated Hydrocarbons: Theory and Crossed Molecular Beams Experiments. *Int. Rev. Phys. Chem.* **2015**, *34*, 461– 514.

(23) Clary, D. C.; Buonomo, E.; Sims, I. R.; Smith, I. W. M.; Geppert, W. D.; Naulin, C.; Costes, M.; Cartechini, L.; Casavecchia, P. C +  $C_2H_2$ : A Key Reaction in Interstellar Chemistry. *J. Phys. Chem.* A **2002**, *106*, 5541–5552.

(24) Kaiser, R. I.; Gu, X. Chemical Dynamics of the Formation of the Ethynylsilylidyne Radical (SiCCH( $X^2\Pi$ )) in the Crossed Beam Reaction of Ground State Silicon Atoms (Si(<sup>3</sup>P)) with Acetylene ( $C_2H_2(X^1\sum_g^+)$ ). J. Chem. Phys. **2009**, 131, No. 104311.

(25) Canosa, A.; Le Picard, S. D.; Gougeon, S.; Rebrion-Rowe, C.; Travers, D.; Rowe, B. R. Rate Coefficients for the Reactions of  $Si({}^{3}P_{J})$  with  $C_{2}H_{2}$  and  $C_{2}H_{4}$ : Experimental Results Down to 15 K. J. Chem. Phys. **2001**, 115, 6495–6503.

(26) Basu, S. C.; Husain, D. Measurement of Absolute Rate Data for the Reactions of Ground State Atomic Silicon  $Si(3^{3}P_{J})$  with Acetylenes by Time-Resolved Atomic Resonance Absorption Spectroscopy. J. Photochem. Photobiol., A **1988**, 42, 1–12.

(27) Basu, S. C.; Husain, D. Absolute Rate Data for the Reaction of Ground State Atomic Silicon,  $Si(3^3P_j)$ , with Olefins Determined by Time-Resolved Atomic Resonance Absorption Spectroscopy. *Z. Phys. Chem.* **1988**, *158*, 1–13.

(28) Talbi, D. An Extensive Ab Initio Study of the Si +  $C_2H_2$  and Si +  $C_2H_4$  Reactions in Relation to the Silicon Astrochemistry. *Chem. Phys.* **2005**, 313, 17–23.

(29) Leonori, F.; Balucani, N.; Capozza, G.; Segoloni, E.; Volpi, G. G.; Casavecchia, P. Dynamics of the  $O({}^{3}P) + C_{2}H_{2}$  Reaction from Crossed Molecular Beam Experiments with Soft Electron Ionization Detection. *Phys. Chem. Chem. Phys.* **2014**, *16*, 10008–10022.

(30) Fu, B.; Han, Y.-C.; Bowman, J. M.; Angelucci, L.; Balucani, N.; Leonori, F.; Casavecchia, P. Intersystem Crossing and Dynamics in  $O(^{3}P) + C_{2}H_{4}$  Multichannel Reaction: Experiment Validates Theory. *Proc. Natl. Acad. Sci. U.S.A.* **2012**, *109*, 9733–9738.

(31) Leonori, F.; Balucani, N.; Nevrly, V.; Bergeat, A.; Falcinelli, S.; Vanuzzo, G.; Casavecchia, P.; Cavallotti, C. Experimental and Theoretical Studies on the Dynamics of the  $O(^{3}P)$  + Propene Reaction: Primary Products, Branching Ratios, and Role of Intersystem Crossing. J. Phys. Chem. C 2015, 119, 14632–14652.

(32) Leonori, F.; Occhiogrosso, A.; Balucani, N.; Bucci, A.; Petrucci, R.; Casavecchia, P. Crossed Molecular Beam Dynamics Studies of the  $O(^{3}P)$  + Allene Reaction: Primary Products, Branching Ratios, and Dominant Role of Intersystem Crossing. *J. Phys. Chem. Lett.* **2012**, *3*, 75–80.

(33) Caracciolo, A.; Vanuzzo, G.; Balucani, N.; Stranges, D.; Casavecchia, P.; Pratali Maffei, L.; Cavallotti, C. Combined Experimental and Theoretical Studies of the  $O({}^{3}P)$  + 1-Butene Reaction Dynamics: Primary Products, Branching Fractions, and Role of Intersystem Crossing. J. Phys. Chem. A **2019**, 123, 9934–9956.

(34) Thomas, A. M.; Dangi, B. B.; Yang, T.; Kaiser, R. I.; Lin, L.; Chou, T.-J.; Chang, A. H. H. Are Nonadiabatic Reaction Dynamics the Key to Novel Organosilicon Molecules? The Silicon  $(Si(^{3}P))$ – Dimethylacetylene  $(C_{4}H_{6}(X^{1}A_{1g}))$  System as a Case Study. J. Phys. Chem. Lett. **2018**, 9, 3340–3347.

(35) Thomas, A. M.; Dangi, B. B.; Yang, T.; Kaiser, R. I.; Sun, B.-J.; Chou, T.-J.; Chang, A. H. H. A Crossed Molecular Beams Investigation of the Reactions of Atomic Silicon (Si(<sup>3</sup>P)) with  $C_4H_6$ Isomers (1,3-Butadiene, 1,2-Butadiene, and 1-Butyne). *Chem. Phys.* **2019**, 520, 70–80. (36) Kaiser, R. I.; Mebel, A. M.; Chang, A. H. H.; Lin, S. H.; Lee, Y. T. Crossed-Beam Reaction of Carbon Atoms with Hydrocarbon Molecules. V. Chemical Dynamics of  $n-C_4H_3$  Formation from Reaction of  $C({}^{3}P_j)$  with Allene,  $H_2CCCH_2$  (X<sup>1</sup>A<sub>1</sub>). *J. Chem. Phys.* **1999**, *110*, 10330–10344.

(37) Stahl, F.; Schleyer, P. vR.; Schaefer, H. F., III; Kaiser, R. I. Reactions of Ethynyl Radicals as a Source of  $C_4$  and  $C_5$  Hydrocarbons in Titan's Atmosphere. *Planet. Space Sci.* **2002**, *50*, 685–692.

(38) Balucani, N.; Mebel, A. M.; Lee, Y. T.; Kaiser, R. I. A Combined Crossed Molecular Beam and Ab Initio Study of the Reactions  $C_2(X^1\sum_g^+, a^3\Pi_u) + C_2H_4 \rightarrow n-C_4H_3(X^2A') + H(^2S_{1/2})$ . J. Phys. Chem. A **2001**, 105, 9813–9818.

(39) Stahl, F.; von Ragué Schleyer, P.; Bettinger, H. F.; Kaiser, R. I.; Lee, Y. T.; Schaefer, H. F., III Reaction of the Ethynyl Radical,  $C_2H$ , with Methylacetylene, CH<sub>3</sub>CCH, Under Single Collision Conditions: Implications for Astrochemistry. *J. Chem. Phys.* **2001**, *114*, 3476– 3487.

(40) Gu, X.; Guo, Y.; Kawamura, E.; Kaiser, R. I. Characteristics and Diagnostics of an Ultrahigh Vacuum Compatible Laser Ablation Source for Crossed Molecular Beam Experiments. *J. Vac. Sci. Technol.,* A **2006**, *24*, 505–511.

(41) Kaiser, R. I.; Chiong, C. C.; Asvany, O.; Lee, Y. T.; Stahl, F.; von R. Schleyer, P.; Schaefer, H. F., III Chemical Dynamics of D1-Methyldiacetylene (CH<sub>3</sub>CCCCD; X<sup>1</sup>A<sub>1</sub>) and D1-Ethynylallene (H<sub>2</sub>CCCH(C<sub>2</sub>D); X<sup>1</sup>A') Formation from Reaction of C<sub>2</sub>D(X<sup>2</sup> $\Sigma^+$ ) with Methylacetylene, CH<sub>3</sub>CCH (X<sup>1</sup>A<sub>1</sub>). J. Chem. Phys. **2001**, 114, 3488–3496.

(42) Guo, Y.; Gu, X.; Kawamura, E.; Kaiser, R. I. Design of a Modular and Versatile Interlock System for Ultrahigh Vacuum Machines: A Crossed Molecular Beam Setup as a Case Study. *Rev. Sci. Instrum.* **2006**, 77, No. 034701.

(43) Yang, T.; Dangi, B. B.; Kaiser, R. I.; Chao, K. H.; Sun, B. J.; Chang, A. H. H.; Nguyen, T. L.; Stanton, J. F. Gas-Phase Formation of the Disilavinylidene (H<sub>2</sub>SiSi) Transient. *Angew. Chem., Int. Ed.* **2017**, *56*, 1264–1268.

(44) Levine, R. D. *Molecular Reaction Dynamics*; Cambridge University Press: Cambridge, U.K., 2005.

(45) Levine, R. D. *Molecular Reaction Dynamics*; Cambridge University Press: Cambridge, 2005.

(46) Miller, W. B.; Safron, S. A.; Herschbach, D. R. Exchange Reactions of Alkali Atoms with Alkali Halides: A Collision Complex Mechanism. *Discuss. Faraday Soc.* **1967**, *44*, 108–122.